

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 209/42	A1	(11) International Publication Number: WO 00/03983 (43) International Publication Date: 27 January 2000 (27.01.00)
(21) International Application Number: PCT/EP99/04943 (22) International Filing Date: 13 July 1999 (13.07.99) (30) Priority Data: 9815483.4 16 July 1998 (16.07.98) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): FEDOULOFF, Michael [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). STRACHAN, John, Bryce [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). (74) Agent: GARRETT, Michael; SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: PROCESS FOR THE PREPARATION OF AN INDOLE DERIVATIVE (57) Abstract <p>A process for the preparation of methyl 3-chloro-3H-indole-3-carboxylate, which comprises reacting an appropriate 3-carboxylate indole compound with N-chlorosuccinimide in the presence of a tertiary amine which is less nucleophilic than DABCO and which has a pK_b of from 8 to 11.</p>		

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR THE PREPARATION OF AN INDOLE DERIVATIVE

This invention relates to a new synthetic process to a compound having
5 pharmacological activity.

WO 93/18036 (SmithKline Beecham plc) describes certain indole compounds
having 5-HT₄ receptor antagonist activity including the compound of formula (I)



(I)


10

and its pharmaceutically acceptable salts. This compound is N-[(1-ⁿbutyl-4-piperidyl)methyl]-3,4-dihydro-2H-[1,3]oxazino[3,2-a]indole-10-carboxamide, referred to herein by its code number SB-207266, (the hydrochloride salt is SB-207266-A), which is being developed by SmithKline Beecham plc as the active
15 ingredient in a medicament for treatment of irritable bowel syndrome.

Example 3 of WO 93/18036 describes a method of preparation of SB-207266-A from N-[(1-ⁿbutyl-4-piperidyl)methyl]indole-3-carboxamide (i.e. the compound corresponding to SB-207266, without the oxazino moiety), by reacting with N-chlorosuccinimide and 3-bromo-1-propanol, followed by treatment with
20 sodium carbonate. N-[(1-ⁿbutyl-4-piperidyl)methyl]indole-3-carboxamide is prepared by coupling N-(1-ⁿbutyl-4-piperidyl)methylamine with an indole-3-carboxylic acid.

WO 98/07728 (SmithKline Beecham plc) describes a process for preparing SB-207266-A which involves the use of the N-(1-ⁿbutyl-4-piperidyl)methylamine intermediate at a later stage in the process thus resulting in an increased yield of SB-207266-A relative to the amount of this intermediate, which is relatively expensive to produce. In particular, the alternative process comprises the reaction of N-(1-ⁿbutyl-4-piperidyl)methylamine with a compound of formula (A):

25




(A)

wherein R is alkyl, such as methyl or ethyl.

- 5 The compound of formula (A) wherein R is methyl is methyl 3,4-dihydro-2H-[1,3]-oxazino[3,2-a]indole-10-carboxylate.

WO98/07728 also describes the preparation of the oxazinoindole compound of the formula (A) from the corresponding indole by reaction with N-chlorosuccinimide and a 3-halo-propanol, such as 3-chloropropanol or 3-bromopropanol followed by
10 cyclisation of the intermediate (B) by treatment with base in a suitable solvent.




(B)

15

The Description in the latter specification describes in more detail the preparation of compound (B) from the corresponding methyl indole-3-carboxylate by reaction of the latter with N-chlorosuccinimide in the presence of 1,4-diazabicyclo[2.2.2]octane (DABCO) to form an intermediate of formula (C):

20




(C)

and subsequent reaction of (C) with 3-chloropropanol in the presence of methane sulphonic acid.

We have now found that the replacement of DABCO by a tertiary amine which is less nucleophilic than DABCO and which has a pK_b of from 8 to 11, especially 1,4-dimethylpiperazine, results in significant advantages for the commercial operation of the above reaction.

According to a feature of the present invention we provide a process for the preparation of the compound of formula (C) above, namely methyl 3-chloro-3H-indole-3-carboxylate, which comprises reacting a compound of formula (D) :



(D)

(in which R is as hereinbefore defined)

with N-chlorosuccinimide in the presence of a tertiary amine which is less nucleophilic than DABCO and which has a pK_b of from 8 to 11, especially 1,4-dimethylpiperazine (DMP).

Other examples of preferred bases for use in accordance with the invention are tetramethylethylenediamine and N-methylpiperidine.

The use of the above amine in place of DABCO has been found to increase significantly the overall yield of the process. The former amine also has the following advantages over DABCO:-

- a) it is non-hygroscopic;
- b) it does not react with dichloromethane, a preferred solvent for the reaction;
- c) the product (C) is more stable allowing extended addition times and better temperature control;
- d) quicker wash separations are possible during commercial manufacture; and
- e) levels of the corresponding 2-methoxy compound, as an impurity, do not increase with extended process times.

The reaction is conveniently effected in an organic solvent such as dichloromethane or chloroform at a temperature in the range -20°C to $+20^{\circ}\text{C}$.

The resulting product of formula (C) can be used for the next step in the synthesis of SB-207266 e.g as described in WO 98/07728 without any need for isolation or purification.

5 The following Example illustrates the invention.


Example

Methyl 2-(3-chloropropoxy)-indole-3-carboxylate (formula (B))

A mixture of methyl indole-3-carboxylate and dichloromethane is cooled to 0° C. 1,4-dimethylpiperazine (0.55eq.) and N-chlorosuccinimide (1.1 eq) are added and the
10 mixture left to stir for two hours to give slurry containing the compound of formula (C) above. The resulting slurry is added to a solution of 3-chloropropanol (1.1 eq) and trichloroacetic acid (0.12 eq) in dichloromethane, maintaining the temperature below 0° C. The reaction mixture is left to stir for half an hour, then washed with 10% aqueous sodium carbonate, 0.5 M hydrochloric acid and water. The organic solution
15 is dried over sodium sulphate, filtered and the solvent evaporated. Toluene is added and the mixture stirred at 0-5° C for one hour. The product is then filtered, washed with toluene and dried to give the title product in 83 % yield.

Claims

1. A process for the preparation of methyl 3-chloro-3H-indole-3-carboxylate,
5 which comprises reacting a compound of formula (D) :



(D)

- (in which R is alkyl)
- 10 with N-chlorosuccinimide in the presence of a tertiary amine which is less nucleophilic than DABCO and which has a pK_b of from 8 to 11.
2. A process as claimed in claim 1 in which the amine is 1,4-dimethylpiperazine.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/04943

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D209/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 07728 A (SMITHKLINE BEECHAM PLC) 26 February 1998 (1998-02-26) cited in the application page 8, line 21 -page 10 -----	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 October 1999

Date of mailing of the international search report

08/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/04943

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9807728 A	26-02-1998	AU 4204897 A	06-03-1998
		CZ 9900516 A	14-07-1999
		EP 0922048 A	16-06-1999
		NO 990697 A	15-02-1999
		PL 331647 A	02-08-1999
